


Claims

We claim:

1. A method for preparation of a compound of the following formula XVI,
5 or salt thereof:



- where
- 10 G is an aryl or heterocyclo group, where said group is mono- or polycyclic, and which is optionally substituted at one or more positions;
Z₁ is O, S, NH, or NR⁶;
Z₂ is O, S, NH, or NR⁶;
A₁ is CR⁷ or N;
 - 15 A₂ is CR⁷ or N;
Y' is J-J'-J'' where J is (CR⁷R^{7'})_n and n = 0-3, J' is O, S, S=O, SO₂, NH, NR⁷,
OP=OOR², OC=O, NR¹C=O, OP=ONHR², OSO₂, NHNH, NHNR⁶, NR⁶NH,
or N=N, and J'' is (CR⁷R^{7'})_n and n = 0-3;
Q₁ is H, alkyl or substituted alkyl, alkenyl or substituted alkenyl, cycloalkyl or
 - 20 substituted cycloalkyl, cycloalkenyl or substituted cycloalkenyl,
heterocycloalkyl or substituted heterocycloalkyl, arylalkyl or substituted
arylalkyl, alkynyl or substituted alkynyl, aryl or substituted aryl, heterocyclo or
substituted heterocyclo, halo, CN, R¹OC=O, R⁴C=O, R⁵R⁶NC=O, HOCR⁷R^{7'},
nitro, R¹OCH₂, R¹O, NH₂, C=OSR¹, SO₂R¹ or NR⁴R⁵;
 - 25 Q₂ is H, alkyl or substituted alkyl, alkenyl or substituted alkenyl, cycloalkyl or
substituted cycloalkyl, cycloalkenyl or substituted cycloalkenyl,
heterocycloalkyl or substituted heterocycloalkyl, arylalkyl or substituted
arylalkyl, alkynyl or substituted alkynyl, aryl or substituted aryl, heterocyclo or

substituted heterocyclo, halo, CN, $R^1OC=O$, $R^4C=O$, $R^5R^6NC=O$, $HO CR^7R^7'$,
nitro, R^1OCH_2 , R^1O , NH_2 , $C=OSR^1$, SO_2R^1 or NR^4R^5 ;


L is a bond, $(CR^7R^7')_n$, NH, NR^5 or $NR^5(CR^7R^7')_n$, where n = 0-3;

- 5 R^1 and $R^{1'}$ are each independently H, alkyl or substituted alkyl, alkenyl or substituted alkenyl, alkynyl or substituted alkynyl, cycloalkyl or substituted cycloalkyl, cycloalkenyl or substituted cycloalkenyl, heterocyclo or substituted heterocyclo, cycloalkylalkyl or substituted cycloalkylalkyl, cycloalkenylalkyl or substituted cycloalkenylalkyl, heterocycloalkyl or substituted heterocycloalkyl, aryl or substituted aryl, arylalkyl or substituted arylalkyl;
- 10 R^2 is alkyl or substituted alkyl, alkenyl or substituted alkenyl, alkynyl or substituted alkynyl, cycloalkyl or substituted cycloalkyl, cycloalkenyl or substituted cycloalkenyl, heterocyclo or substituted heterocyclo, cycloalkylalkyl or substituted cycloalkylalkyl, cycloalkenylalkyl or substituted cycloalkenylalkyl, heterocycloalkyl or substituted heterocycloalkyl, aryl or substituted aryl, arylalkyl or substituted arylalkyl;
- 15 R^4 is H, alkyl or substituted alkyl, alkenyl or substituted alkenyl, alkynyl or substituted alkynyl, cycloalkyl or substituted cycloalkyl, cycloalkenyl or substituted cycloalkenyl, heterocyclo or substituted heterocyclo, cycloalkylalkyl or substituted cycloalkylalkyl, cycloalkenylalkyl or substituted cycloalkenylalkyl, heterocycloalkyl or substituted heterocycloalkyl, aryl or substituted aryl, arylalkyl or substituted arylalkyl, $R^1C=O$, $R^1NHC=O$, SO_2OR^1 , or $SO_2NR^1R^{1'}$;
- 20 R^5 is alkyl or substituted alkyl, alkenyl or substituted alkenyl, alkynyl or substituted alkynyl, cycloalkyl or substituted cycloalkyl, cycloalkenyl or substituted cycloalkenyl, heterocyclo or substituted heterocyclo, cycloalkylalkyl or substituted cycloalkylalkyl, cycloalkenylalkyl or substituted cycloalkenylalkyl, heterocycloalkyl or substituted heterocycloalkyl, aryl or substituted aryl, arylalkyl or substituted arylalkyl, $R^1C=O$, $R^1NHC=O$, SO_2R^1 , SO_2OR^1 , or $SO_2NR^1R^{1'}$;
- 25 R^6 is alkyl or substituted alkyl, alkenyl or substituted alkenyl, alkynyl or substituted alkynyl, cycloalkyl or substituted cycloalkyl, cycloalkenyl or substituted cycloalkenyl, heterocyclo or substituted heterocyclo, cycloalkylalkyl or substituted cycloalkylalkyl, cycloalkenylalkyl or substituted cycloalkenylalkyl, heterocycloalkyl or substituted heterocycloalkyl, aryl or substituted aryl, arylalkyl or substituted arylalkyl, $R^1C=O$, $R^1NHC=O$, SO_2R^1 , SO_2OR^1 , or $SO_2NR^1R^{1'}$;
- 30 R^6 is alkyl or substituted alkyl, alkenyl or substituted alkenyl, alkynyl or substituted alkynyl, cycloalkyl or substituted cycloalkyl, cycloalkenyl or substituted cycloalkenyl, heterocyclo or substituted heterocyclo, cycloalkylalkyl or substituted cycloalkylalkyl, cycloalkenylalkyl or substituted cycloalkenylalkyl, heterocycloalkyl or substituted heterocycloalkyl, aryl or substituted aryl, arylalkyl or substituted arylalkyl, $R^1C=O$, $R^1NHC=O$, SO_2R^1 , SO_2OR^1 , or $SO_2NR^1R^{1'}$;

cycloalkenyl, heterocyclo or substituted heterocyclo, cycloalkylalkyl or substituted cycloalkylalkyl, cycloalkenylalkyl or substituted cycloalkenylalkyl, heterocycloalkyl or substituted heterocycloalkyl, aryl or substituted aryl, arylalkyl or substituted arylalkyl, CN, OH, OR¹, R¹C=O, R¹NHC=O, SO₂R¹, SO₂OR¹, or SO₂NR¹R^{1'}; and

R⁷ and R^{7'} are each independently H, alkyl or substituted alkyl, alkenyl or substituted alkenyl, alkynyl or substituted alkynyl, cycloalkyl or substituted cycloalkyl, cycloalkenyl or substituted cycloalkenyl, heterocyclo or substituted heterocyclo, cycloalkylalkyl or substituted cycloalkylalkyl, cycloalkenylalkyl or substituted cycloalkenylalkyl, heterocycloalkyl or substituted heterocycloalkyl, aryl or substituted aryl, arylalkyl or substituted arylalkyl, halo, CN, OR¹, nitro, hydroxylamine, hydroxylamide, amino, NHR⁴, NR²R⁵, NOR¹, thiol, alkylthio or substituted alkylthio, R¹C=O, R¹(C=O)O, R¹OC=O, R¹NHC=O, SO₂R¹, SOR¹, PO₃R¹R^{1'}, R¹R^{1'}NC=O, C=OSR¹, SO₂R¹, SO₂OR¹, or SO₂NR¹R^{1'};

comprising the steps of contacting a compound of the following formula XV, or salt thereof:




20

where the symbols are as defined above;

with an enzyme or microorganism capable of catalyzing the hydroxylation of said compound XV to said compound XVI, and effecting said hydroxylation.

25

2. A method for preparation of a compound of the following formula XVIII, or salt thereof:



where

G is an aryl or heterocyclo group, where said group is mono- or polycyclic, and which is optionally substituted at one or more positions;

5 Z_1 is O, S, NH, or NR^6 ;

Z_2 is O, S, NH, or NR^6 ;

A_1 is CR^7 or N;

A_2 is CR^7 or N;

Y' is J-J'-J'' where J is $(CR^7R^{7'})_n$ and $n = 0-3$, J' is O, S, S=O, SO_2 , NH, NR^7 ,

10 $OP=OOR^2$, $OC=O$, $NR^1C=O$, $OP=ONHR^2$, OSO_2 , $NHNH$, $NHNR^6$, NR^6NH , or $N=N$, and J'' is $(CR^7R^{7'})_n$ and $n = 0-3$;

Q_1 is H, alkyl or substituted alkyl, alkenyl or substituted alkenyl, cycloalkyl or

substituted cycloalkyl, cycloalkenyl or substituted cycloalkenyl,

heterocycloalkyl or substituted heterocycloalkyl, arylalkyl or substituted

15 arylalkyl, alkynyl or substituted alkynyl, aryl or substituted aryl, heterocyclo or substituted heterocyclo, halo, CN, $R^1OC=O$, $R^4C=O$, $R^5R^6NC=O$, $HO CR^7R^{7'}$, nitro, R^1OCH_2 , R^1O , NH_2 , $C=OSR^1$, SO_2R^1 or NR^4R^5 ;

Q_2 is H, alkyl or substituted alkyl, alkenyl or substituted alkenyl, cycloalkyl or

substituted cycloalkyl, cycloalkenyl or substituted cycloalkenyl,

20 heterocycloalkyl or substituted heterocycloalkyl, arylalkyl or substituted

arylalkyl, alkynyl or substituted alkynyl, aryl or substituted aryl, heterocyclo or

substituted heterocyclo, halo, CN, $R^1OC=O$, $R^4C=O$, $R^5R^6NC=O$, $HO CR^7R^{7'}$,

nitro, R^1OCH_2 , R^1O , NH_2 , $C=OSR^1$, SO_2R^1 or NR^4R^5 ;

L is a bond, $(CR^7R^{7'})_n$, NH, NR^5 or $NR^5(CR^7R^{7'})_n$, where $n = 0-3$;

25 R^1 and $R^{1'}$ are each independently H, alkyl or substituted alkyl, alkenyl or substituted alkenyl, alkynyl or substituted alkynyl, cycloalkyl or substituted cycloalkyl,

cycloalkenyl or substituted cycloalkenyl, heterocyclo or substituted

heterocyclo, cycloalkylalkyl or substituted cycloalkylalkyl, cycloalkenylalkyl or

substituted cycloalkenylalkyl, heterocycloalkyl or substituted heterocycloalkyl,

30 aryl or substituted aryl, arylalkyl or substituted arylalkyl;

R^2 is alkyl or substituted alkyl, alkenyl or substituted alkenyl, alkynyl or substituted

alkynyl, cycloalkyl or substituted cycloalkyl, cycloalkenyl or substituted

cycloalkenyl, heterocyclo or substituted heterocyclo, cycloalkylalkyl or substituted cycloalkylalkyl, cycloalkenylalkyl or substituted cycloalkenylalkyl, heterocycloalkyl or substituted heterocycloalkyl, aryl or substituted aryl, arylalkyl or substituted arylalkyl;

- 5 R^4 is H, alkyl or substituted alkyl, alkenyl or substituted alkenyl, alkynyl or substituted alkynyl, cycloalkyl or substituted cycloalkyl, cycloalkenyl or substituted cycloalkenyl, heterocyclo or substituted heterocyclo, cycloalkylalkyl or substituted cycloalkylalkyl, cycloalkenylalkyl or substituted cycloalkenylalkyl, heterocycloalkyl or substituted heterocycloalkyl, aryl or substituted aryl, arylalkyl or substituted arylalkyl, $R^1C=O$, $R^1NHC=O$, SO_2OR^1 , or $SO_2NR^1R^{1'}$;


- 10 R^5 is alkyl or substituted alkyl, alkenyl or substituted alkenyl, alkynyl or substituted alkynyl, cycloalkyl or substituted cycloalkyl, cycloalkenyl or substituted cycloalkenyl, heterocyclo or substituted heterocyclo, cycloalkylalkyl or substituted cycloalkylalkyl, cycloalkenylalkyl or substituted cycloalkenylalkyl, heterocycloalkyl or substituted heterocycloalkyl, aryl or substituted aryl, arylalkyl or substituted arylalkyl, $R^1C=O$, $R^1NHC=O$, SO_2R^1 , SO_2OR^1 , or $SO_2NR^1R^{1'}$;

- 15 R^6 is alkyl or substituted alkyl, alkenyl or substituted alkenyl, alkynyl or substituted alkynyl, cycloalkyl or substituted cycloalkyl, cycloalkenyl or substituted cycloalkenyl, heterocyclo or substituted heterocyclo, cycloalkylalkyl or substituted cycloalkylalkyl, cycloalkenylalkyl or substituted cycloalkenylalkyl, heterocycloalkyl or substituted heterocycloalkyl, aryl or substituted aryl, arylalkyl or substituted arylalkyl, CN, OH, OR^1 , $R^1C=O$, $R^1NHC=O$, SO_2R^1 , SO_2OR^1 , or $SO_2NR^1R^{1'}$; and

- 20 R^7 and $R^{7'}$ are each independently H, alkyl or substituted alkyl, alkenyl or substituted alkenyl, alkynyl or substituted alkynyl, cycloalkyl or substituted cycloalkyl, cycloalkenyl or substituted cycloalkenyl, heterocyclo or substituted heterocyclo, cycloalkylalkyl or substituted cycloalkylalkyl, cycloalkenylalkyl or substituted cycloalkenylalkyl, heterocycloalkyl or substituted heterocycloalkyl, aryl or substituted aryl, arylalkyl or substituted arylalkyl,
- 25
- 30

halo, CN, OR¹, nitro, hydroxylamine, hydroxylamide, amino, NHR⁴, NR²R⁵, NOR¹, thiol, alkylthio or substituted alkylthio, R¹C=O, R¹(C=O)O, R¹OC=O, R¹NHC=O, SO₂R¹, SOR¹, PO₃R¹R^{1'}, R¹R^{1'}NC=O, C=OSR¹, SO₂R¹, SO₂OR¹, or SO₂NR¹R^{1'};

- 5 comprising the steps of contacting a compound of the following formula XVII, or salt thereof:



XVII

- where the symbols are as defined above;
- 10 with an enzyme or microorganism capable of catalyzing the opening of the epoxide ring of compound XVII to form the diol of said compound XVIII, and effecting said ring opening and diol formation.